

ANTIDIABETIC ACTIVITY OF ETHANOLIC EXTRACT OF *Kalanchoe pinnata* LEAVES IN ALLOXAN INDUCED HYPERGLYCAEMIC RATS

Tri Yuliani, Indah D. Dewijanti, Sofna D.S. Banjarnahor

Research Center for
Chemistry, Indonesian
Institute of Sciences (LIPI),
Kawasan PUSPITEK, Muncul,
Serpong, Kota Tangerang
Selatan, Banten 15314

Submitted: 13-5-2016

Revised: 25-07-2016

Accepted: 11-09-2016

*Corresponding author
Tri Yuliani

Email:
anik_baik@yahoo.com

ABSTRACT

Diabetes remains a major burden in health care both in developed and developing countries. *Kalanchoe pinnata* has been used as a traditional medicine to treat diabetes. The aim of this study to find scientific evidence of antidiabetic activity of *Kalanchoe pinnata* extract (KPE) through hypoglycemic effect using animal model of diabetes mellitus. Hyperglycaemia was developed in rats using alloxan 150mg/kgBW. Three days after alloxan injection, rats having fasting blood glucose (FBG) >200mg/dL were divided into six groups, namely HG (hyperglycaemia), HG+KPE high-dose (hyperglycaemia+KPE 33.2mg/kg), HG+KPE medium-dose (hyperglycaemia+KPE 11.6mg/kg), HG+KPE low-dose (hyperglycaemia+KPE 5.8mg/kg), standard drug 1 (hyperglycaemia+glibenclamide 1.35mg/kg), standard drug 2 (hyperglycaemia+acarbose 13.5mg/kg). Then, FBG was measured every 5 days recorded as t1, t2, and t3 to determine fluctuations in blood glucose. At the end of the study, rats were sacrificed, pancreas was collected and number of pancreatic beta cell langerhans was determined. KPE 11.6mg/kg showed best hypoglycemic effect and improvement of the number of pancreatic beta cell langerhans. KPE has hypoglycemic effect through improvement of the number of pancreatic beta cell langerhans but not in dose dependent manner.

Key words: Antidiabetic, *Kalanchoe pinnata*, hyperglycaemic, alloxan, pancreatic

INTRODUCTION

WHO (2015) reported 9% people aged 18+ worldwide suffered from diabetes, while in Indonesia the prevalence was only 1.5% (Kemenkes RI, 2013). It is worth noticing that this disease is a burden of health care in both developed and developing countries (Abdulazeez *et al.*, 2013). Diabetes develops from prolong glucose imbalance between intracellular and extracellular resulted in hyperglycaemia state in which blood glucose can not be utilized by living cells. These conditions lead to vascular complication through increasing oxidative stress and inflammation. Vascular complication is the leading cause of death in both diabetes type 1 and type 2 patient (Domingueti *et al.*, 2015).

Diabetes mellitus (DM) are classified as two major disease: type 1 DM and type 2 DM. Type 1 DM is an autoimmune disease that immune system of the body attack insulin producing pancreatic β cells resulting in low

insulin production. This condition leads to insulin dependent DM. In the other hand, type 2 DM was caused by inadequate action of insulin in action site (Hossain *et al.*, 2016).

Pharmacotherapy in DM patient does not eliminate the cause of the disease but its targets on the symptom using insulin and hypoglycemic agent such as sulfonylurea and biguanids (Abdulazeez *et al.*, 2013; Verma *et al.*, 2015). However, sulfonylurea exert several adverse effect such as hypoglycemia and weight gain, in addition to toxic effect on liver and renal, while metformin cause lactic acidosis (Verma *et al.*, 2015).

Alloxan is widely used in experimental pharmacology to induce diabetes in rat. The pathological mechanism is by destruction of insulin producing pancreatic β -cells. Alloxan undergo redox reaction resulted in increase increasing of radical species. In this case, pancreatic β -cells is the most sensitive organ

towards these radical attacks. Hyperglycaemia state does not occur immediately. At first, alloxan induces insulin release independent of glucose level. However, this effect is followed by suppression of the islet response to even high level of glucose leading to hyperglycaemia state (Szkudelski *et al.*, 2001).

Nowadays, the need for safe and effective drugs has brought natural product as the promising source of companion drug for diabetes (Sharma and Gupta, 2015). *Kalanchoe pinnata* has been traditionally used worldwide as medicinal plants, including to treat diabetes (Patil *et al.*, 2013). Our previous study reveals its antidiabetic activity through inhibition of α -glucosidase with $IC_{50} \sim 16.12$ ppm. Meanwhile, radical scavenger activity using DPPH method revealed $IC_{50} \sim 29.61$ ppm of ethanol fraction, 23.15 ppm of butanol fraction, and 17.27 ppm of ethyl acetic fraction (Indah *et al.*, 2013).

Therefore, current study aims to determine antidiabetic activity of ethanolic extract of *Kalanchoe pinnata* in alloxan-induced diabetic rats.

MATERIALS AND METHODS

Kalanchoe pinnata leaves were collected from Kawasan Puspitpek Serpong and were authenticated by Research Center for Biology, Indonesian Institute of Sciences (LIPI).

Preparation of plant extracts

The leaves were dried using blower oven $< 50^{\circ}C$ for 24h, crushed, and extracted using 70% ethanol (3x24h maceration). Ethanol extracts obtained were then concentrated in rotary evaporator under vacuum. The yield of ethanolic extract of *Kalanchoe pinnata* (KP) dry leaves was 15.7%.

Animals

The Male Sprague Dawley rats (150-350g) were procured from Fakultas Kedokteran Hewan IPB, Bogor and housed under standard conditions of temperature and relative humidity with 12h light/dark cycle. Animals were fed on standard commercial pellet diet and water *ad libitum*. The ethical clearance of the experiment has been approved by Health Research Ethics Committee, Universitas Indonesia and Cipto Mangunkusumo Hospital, Indonesia (Ethical

clearance certificate number: 663/UN2.F1/ETIK/2016).

Induction of diabetes

Induction of diabetes was performed according to Misra and Aiman (2012), with modification. Base line FBG was measured before alloxan injection (t(-1)). Alloxan monohydrate (Sigma Aldrich; stored at $4^{\circ}C$) was dissolved in normal saline at room temperature (freshly prepared) and injected to 30 overnight fasted male Sprague-Dawley rats at a dose of 150mg/kg intraperitoneally. The animals were then kept for the next 7 days on 10% glucose after alloxan administration. After 72h of alloxan injection (t0), FBG was determined using GlucoDr glucometer strips. Animals with FBG > 200 mg/dL were considered to have developed experimental diabetes as shown by hyperglycaemia state. Then, FBG was measured every 5 days recorded as t1, t2, and t3 to determine fluctuations in FBG. Blood was collected from tail vein. As standard reference, Glibenclamide was given at a dose of 1.35mg/kg orally per day, while Acarbose was given at a dose of 13.5mg/kg orally per day.

Animal experimentation

In the present study the animals were distributed into 7 groups (n=5) as follows: normal control, HG (hyperglycaemia), HG+KPE high-dose (hyperglycaemia+KPE 33.2mg/kg), HG+KPE medium-dose (hyperglycaemia+KPE 11.6mg/kg), HG+KPE low-dose (hyperglycaemia+KPE 5.8mg/kg), standard drug 1 (hyperglycaemia+ glibenclamide 1.35mg/kg), standard drug 2 (hyperglycaemia+acarbose 13.5mg/kg). The study was conducted for 15 days to evaluate the potential of the extracts to lower FBG level. Body weights of the rats were monitored weekly during the study period.

Pancreas histological observation

Pancreas histological observation was conducted on day 15, at the end of the experimental period. Rats were euthanized by ether and the pancreas were collected and fixed with Bouin's solution for 24h. The pancreas was then stained with *Hematoxylin eosin* (HE) for observation of number of beta cell Langerhans.

Tabel I. Effect of KPE on FBG during study

Group	FBG± SD (mg/L)				
	t (-1)	To	t1	t2	t3
Normal	103.67±15.33	117.92±8.07	107.83±7.15	97.42±13.75	93.5±8.24
HG	115±2.45	429±106.88	267±106.10	207±104.24	130±16.85
HG+KPE (high dose)	112.2±12.76	384±137.48	174.6±111.84	111.2±24.22	116±13.93
HG+KPE (medium dose)	112±10.05	259.4±48.53	134.4±30.00	115.8±17.88	116.2±10.40
HG+KPE (low dose)	112.2±6.57	353.8±163.69	142.4±35.67	113.6±16.36	124.6±10.26
HG+ Acarbose	109.5±16.99	262±36.79	121.83±22.25	105.67±14.51	97±8.65

Statistical analysis

Numeric data were expressed as mean ± SD. Data were analyzed using one-way analysis of variance (ANOVA) ($p < 0.05$) followed by Multiple LSD.

RESULTS DISCUSSION

Base line FBG (t(-1)) was determined before alloxan injection. All rats showed normal FBG with no significant difference among the groups (Table I, Figure 1).

Three days after alloxan injection (t0), hyperglycaemia state was confirmed by FBG > 200 mg/dL in all the rats. However, FBG increase was not significant in HG+KPE medium dose group and acarbose group compared to normal group. Meanwhile, no significant difference was observed among groups that received alloxan injection.

FBG reduction were detected after 5 days (t1) alloxan injection in all groups. While 5 days treatment brought FBG to a level < 200 mg/dL in all groups, HG groups remained suffered from hyperglycaemia as shown by FBG > 200 mg/dL. As a result, only HG group differed significantly from normal group. In the other hand, all treatment reduced FBG significantly, but not high-dose KPE.

Ten days after treatment (t2), all rats' FBG kept decreasing. However, HG group remained in hyperglycaemia state as shown by FBG > 200 mg/dL. As a result, only HG group differed significantly from normal group and all treatments reduced FBG significantly.

Fifteen days after treatment, all rats' FBG went back to normal as shown by FBG < 200 mg/dL. However, only standard

treatment (glibenclamide or acarbose) reduced FBG similar to normal while KPE treatment did not reduced FBG significantly.

Histological observation using HE staining showed that all treatment, but KPE high-dose, improved the morphology of β cells in comparison to the HG group based on the number of cell (Figure 2). However, only KPE medium and low dose increased the number of β cells significantly.

K. pinnata has traditionally been used as medicinal plant to treat various ailments around the world, including diabetes (Patil *et al.*, 2013). Therefore, several study have been conducted to evaluate its bioactivity, such as antidiabetics, anticonvulsant, antinociceptive, antiedematogenic, antiinflammatory, anticancer and antiHPV, antileishmanial (Patil *et al.*, 2013; Mora-Perez *et al.*, 2016; Ferreira *et al.*, 2014; Mahata *et al.*, 2012; Muzitano *et al.*, 2006).

K. pinnata leaves contain anthocyanins that considered responsible for its antidiabetic, anticancer, cardiovascular, and neurological activity (Cruz *et al.*, 2012). Anthocyanins are water soluble flavonoid compounds that gives color in plant (Cruz *et al.*, 2012). Phytochemistry study of the KPE used in this study revealed flavonoids and steroids compound and isolated quercetin glycosides (Fajriyah, 2011).

We suggest that the bioactivity of *K. pinnata* in reducing FBG and improving morphology of the islets of Langerhans and β cells were caused by its various natural compounds constituent, for example quercetin through antioxidant mechanism (Figure 3).




Figure 1. Effect of KPE treatment on FBG level in alloxan induced rat. HG=Alloxan, KPE= *Kalanchoe pinnata* extract, high dose = 33.2mg/kg BW, medium dose= 11.6mg/kg BW, low dose= 5.8mg/kg BW, Glibenclamide = 1.35mg/kg bw, Acarbose = 13.5mg/kg BW. t(-1) =base line, t 0= after alloxan injection, t1 or t2 or t3 = 5 or 10 or 15 days after test sample administration. Data are expressed as mean \pm SD. * & \times , $p < 0.05$ (* indicates comparison between normal and HG; \times indicates comparison between HG and treatments).




Figure 2. Effect of KPE treatment on number of pancreatic beta cell langerhans in alloxan induced rat. HG=Alloxan, KPE= *Kalanchoe pinnata* extract, high dose = 33.2 mg/kgbw, medium dose= 11.6 mg/kg bw, low dose= 5.8 mg/kg bw, Glibenclamide = 1.35 mg/kg bw, Acarbose = 13.5 mg/kg bw. \times , $p < 0.05$. (\times indicates comparison between HG and treatments).




Figure 3. Pancreatic β cell Langerhans stained with *Hematoxylin eosin* (HE).

It is worth noting that oxidative stress plays a role in the pathogenesis of diabetes mellitus. Hyperglycaemia was characterized with increase oxidative stress leading to defect in insulin action and insulin secretion. Quercetin as antioxidants reduced oxidative stress leading to protection of β cells of the pancreas resulted in the increase of insulin production and decreased of FBG. Several studies reported quercetin mechanism of action in diabetes such as decreases lipid peroxidation, increases antioxidant enzymes activity, inhibits insulin-dependent activation of phosphoinositol-3-kinase (PI-3K), and reduces intestinal glucose absorption by inhibiting GLUT (Sunarwidhi *et al.*, 2014).

Based on pharmacological and histopathological studies, we suggested that the hypoglycemic effect of KPE was optimum at medium dose (11.6mg/kg bw).

However, it is difficult to conclude that hypoglycemic activity of KPE in this study is only caused by quercetin. Various active compounds in KPE may synergistically increase hypoglycemic effect.

Therefore, the mechanism of action of each active compounds in the extract needs further investigation, including immuno histochemical observation on pancreatic insulin expression.

CONCLUSION

This study showed that KPE possessed bioactivity in lowering blood glucose and improving the morphology of β cells of Langerhans islet. KPE is potential to be developed as a blood glucose-lowering agent for diabetic patients.

ACKNOWLEDGMENTS

We thank the "Program Insentif Riset Peneliti dan Perekayasa Tahun 2011" for financial support in the study.

REFERENCES

- Abdulazeez SS. 2013. Diabetes treatment: A rapid review of the current and future scope of stem cell research. *Saudi Pharm. J.* 23: 333–340.
- Cruz BP., Chedier LM., Peixoto PHP., Fabri RL., Pimenta DS. 2012. Effects of light intensity on the distribution of anthocyanins in *Kalanchoe brasiliensis* Camb. and *Kalanchoe pinnata* (Lamk.) Pers, *Anais da Academia Brasileira de Ciências.* 84(1): 211-217
- Domingueti CP., Dusse LM., Carvalho MD., de Sousa LP., Gomes KB., Fernandes AP. 2015. Diabetes mellitus: The linkage between oxidative stress, inflammation, hypercoagulability and vascular complications. *J Diabetes Complications.* Dec 18.
- Ferreira RT., Coutinho MAS., Carmo Malvar D., Costa EA., Florentino IF., Costa SS., Vanderlinde FA. 2014. Mechanisms Underlying the Antinociceptive, Antiedematogenic, and Anti-Inflammatory Activity of the Main Flavonoid from *Kalanchoe pinnata*. *Evidence-Based Complementary and Alternative Medicine:* 429256.
- Global status report on noncommunicable diseases. 2014. Geneva, World Health Organization.
- Hossain MK., Dayem AA., Han J., Saha SK., Yang G., Choi HY., Cho S. 2016. Recent Advances in Disease Modeling and Drug Discovery for Diabetes Mellitus Using Induced Pluripotent Stem Cells. *Int. J. Mol. Sci.* 17: 256.
- Indah DD., Euis F, Megawati, Tri Y. 2012. The Antidiabetic Activity of Cocor Bebek Leaves' (*Kalanchoe pinnata* Lam.Pers.) Ethanolic Extract from Various Areas. *J.Trop. Life Sciences.* 2(2) : 37 – 39.
- Kementrian Kesehatan RI. Badan Penelitian dan Pengembangan. Riskesdas. 2013. Jakarta.
- Mahata S., Maru S., Shukla S., Pandey A., Mughesh G., Das BC., Bharti AC. 2012. Anticancer property of Bryophyllum pinnata (Lam.) Oken. leaf on human cervical cancer cells. *BMC Complementary and Alternative Medicine.* 12:15
- Misra M. and Aiman U. 2012. Alloxan: An unpredictable drug for diabetes induction? *Indian J Pharmacol.*44(4): 538–539.
- Muzitano MF, Tinoco LW, Guette C, Kaiser CR, RossiBergmann B, Costa SS. The antileishmanial activity assessment of unusual flavonoids from *Kalanchoe*

- pinnata. *Phytochemistry*, 2006; 67: 2071-2077
- Patil SB., Dongare VR., Kulkarni CR., Joglekar MM., Arvindekar AU. 2013 Antidiabetic activity of *Kalanchoe pinnata* in streptozotocin-induced diabetic rats by glucose independent insulin secretagogue action. *Pharm Biol.* 51(11):1411-8.
- Sharma SB., Gupta R. 2015. Drug development from natural resource: a systematic approach. *Mini Rev Med Chem.* 15(1):52-7.
- Sunarwidhi AL., Sudarsono S., Nugroho AE. 2014. Hypoglycemic Effect of Combination of *Azadirachta indica* A. Juss. and *Gynura procumbens* (Lour.) Merr. Ethanolic Extracts Standardized by Rutin and Quercetin in Alloxan-induced Hyperglycaemia Rats. *Adv Pharm Bull.* 4(Suppl 2): 613-618.
- Szkudelski T. 2001. The Mechanism of Alloxan and Streptozotocin Action in B Cells of the Rat Pancreas. *Physiol. Res.* 50: 536-546.
- Verma RK., Mishra G., Singh P., Jha KK., Khosa RL. 2015. Anti-diabetic activity of methanolic extract of *Alpinia galanga* Linn. aerial parts in streptozotocin induced diabetic rats. *Ayu.* 36(1):91-95.